

Dicke-Type Energy Level Crossings in Cavity-Induced Atom Cooling: Another Superradiant Cooling

Masao Hirokawa^{1,*}

¹*Department of Mathematics, Okayama University, Okayama 700-8530, Japan*

(Dated: December 19, 2008)

This paper is devoted to energy-spectral analysis for the system of a two-level atom coupled with photons in a cavity. It is shown that the Dicke-type energy level crossings take place when the atom-cavity interaction of the system undergoes changes between the weak coupling regime and the strong one. Using the phenomenon of the crossings we develop the idea of cavity-induced atom cooling proposed by the group of Ritsch, and we lay mathematical foundations of a possible mechanism for another superradiant cooling in addition to that proposed by Domokos and Ritsch. The process of our superradiant cooling can function well by cavity decay and by control of the position of the atom, at least in (mathematical) theory, even if there is neither atomic absorption nor atomic emission of photons.

PACS numbers: 37.10.De, 42.50.Pq, 02.30.Sa, 02.30.Tb

I. INTRODUCTION

Laser cooling is one of attractive subjects of modern physics. It has been demonstrated with several experimental techniques such as the ion cooling [1], the Doppler cooling [2], the Sisyphus cooling [3], etc. Also it has enabled us to observe many fundamental phenomena in theoretical physics. One of typical instances of applications of the laser cooling is the observation of Bose-Einstein condensation [4, 5]. It has been about ten years since another type of laser cooling was proposed using the strong atom-photon interaction. Such a strong interaction between atom and photons is realized in the so-called cavity quantum electrodynamics (QED) [6, 7, 8, 9, 10, 11]. Thus, the group of Ritsch has investigated the system of a two-level atom coupled with a laser, and then, they have found a mechanism for cooling the atom [12], which is similar to that of the Sisyphus cooling. The cooling mechanism is called *cavity-induced atom cooling*. In the process of their cavity-induced atom cooling, the method to carry away the energy from the atom coupled with photons is given not only by atomic decay (i.e., atomic spontaneous emission of photons) but also by cavity decay. It has experimentally been confirmed that the cavity decay works in the cooling system [13, 14]. Concerning the cooling methods using cavity QED, Domokos and Ritsch proposed a concept of superradiant cooling [15] based on the atomic self-organization and cooperation among many atoms in a cavity [16]. The atom-photon interaction in the strong coupling regime brings the situation amazingly different from ordinary atomic decay [10]. We will adopt this difference into our arguments on the cavity-induced atom cooling.

As well as the ensemble of many two-level atoms coupled with a laser has the possibility of making super-

radiance as a cooperative effect in optics [17, 18], another superradiance may also appear in energy spectrum even for the system of a two-level atom coupled with a laser, provided that it is in the strong coupling regime [19, 20, 21, 22, 23]. As far as atom-laser interaction in the cooling process for the Bose-Einstein condensation goes, a superradiance has been experimentally observed under a certain physical condition [24, 25, 26, 27], and this phenomenon has been theoretically shown [28, 29, 30]. It was pointed out that there is a possibility that superradiance causes the energy level crossing between the initial ground state energy and an initial excited state energy [19, 22, 23], namely, a kind of phase transition occurs. This is an optical phenomenon of light-induced phase transitions though it is not a cooperative effect in optics. The details of such an energy level crossing has precisely been studied, and then, this type of crossing is called the *Dicke-type (energy level) crossing* [31]. We will strictly define its meaning in Sec.II. The reason why we call the energy level crossing so is that it is basically caused by the mathematical mechanism [19, 23] of Dicke's superradiance [32].

This paper is devoted to developing the cavity-induced atom cooling. Namely, we will show that the Dicke-type energy level crossings take place when the system undergoes changes between the weak coupling regime and the strong one. Using the crossings, we will propose a possibility of another superradiant cooling in terms of the energy spectrum from our point of view. In our proposal we will consider whether the followings are possible in theory for cooling the atom in a cavity: 1) can we use a laser only for controlling the strength of the atom-cavity interaction without throwing another laser to the atom for driving it to an excited state? 2) can we expect that the energy loss caused by cavity decay becomes much larger? To perform our research into the problems, we consider an ideal situation only to see the energy-spectral property for our system without considering, for example, the laser heating processes caused by diffusion of the atomic momentum. Some mathematical technique to make the

*Electronic address: hirokawa@math.okayama-u.ac.jp;
URL: <http://www.math.okayama-u.ac.jp/~hirokawa>

spectral analysis for such systems have been developed lately [31, 33, 34, 35, 36]. Thus, another purpose of this paper is to show the mechanism of the Dicke-type energy level crossing in the cavity-induced atom cooling as rigorously as in the works in Ref.37 so that the process of our superradiant cooling can function well in (mathematical) theory.

Our paper is constructed in the following. In Sec.II we will generalize the Hamiltonian $H(\Omega, \alpha; d)$ which Ritsch's group handled, where Ω is a function of space-time point and governs the atom-photon interaction, α also a function of space-time and a generalization of the strength of the pump field, and d a parameter for non-linear coupling of the atom and photons. Moreover, we will define some notion to explain what the Dicke-type energy level crossings take place for $H(\Omega, \alpha; d)$ with $\alpha \equiv 0$. In Sec.IV we will show the existence of the superradiant ground state energy for $H(\Omega, \alpha; d)$ with $\alpha \equiv 0$ and $d = 1$ in the strong coupling regime. In Sec.V we will argue the stability of the Dicke-type energy level crossing under the condition $\alpha \neq 0$.

II. HAMILTONIAN AND SOME NOTION

In Ref.12 the group of Ritsch studied a Hamiltonian adopting dipole and rotating wave approximation. To write down their Hamiltonian, we define some operators: the atomic position (resp. momentum) operator is denoted by x (resp. p), the photon annihilation (resp. creation) operator by a (resp. $a^\dagger$), and the atomic operator is given by $\sigma_{ij} = |i\rangle\langle j|$, $i, j = 0, 1$. Then, the Hamiltonian is

$$H = \frac{1}{2m}p^2 - \Delta\sigma_{11} - \Delta_c a^\dagger a + i\Omega(x)(\sigma_{01}a^\dagger - \sigma_{10}a) + i\alpha(a - a^\dagger),$$

where two real numbers Δ and Δ_c with $-\infty < \Delta < +\infty$ and $\Delta_c < 0$ are respectively the atom-pump detuning and the detuning of the empty cavity relative to the pump frequency, and $\Omega(x)$ stands for the atom-cavity coupling constant, i.e., $\Omega(x) = \Omega_0 \cos kx$ with the position x of the atom and the wave number k of photons of the laser. We note that in the case $\alpha = 0$ the Hamiltonian H is used to argue the resonant interaction of an atom with a microwave field [38]. In Hamiltonian H , the part consisting of the first, the second, and the third terms (i.e., $(2m)^{-1}p^2 - \Delta\sigma_{11} - \Delta_c a^\dagger a$) is the free Hamiltonian of our system. Each of the fourth and fifth terms represents the Hamiltonian of interaction and the energy operator of the pump field respectively. In this section we generalize the above Hamiltonian. Our generalization is the following: (1) we consider not only the linear coupling but also non-linear coupling; (2) we introduce the time-dependence into the coupling constant $\Omega(x)$; (3) we con-

sider the general operator which represents not only the energy operator of the pump field but also, for instance, the energy operator of the pump field plus some error potential coming from the environment of the experiment for testing the system. Thus our Hamiltonian reads

$$H(\Omega, \alpha; d) = \frac{1}{2m}p^2 - \Delta\sigma_{11} - \Delta_c a^\dagger a + i\Omega(x, t)(\sigma_{01}a^{\dagger d} - \sigma_{10}a^d) + \alpha(x, t)W(x, t)$$

for $d = 1, 2, \dots$, where $\Omega(x, t)$ and $\alpha(x, t)$ are continuous, real-valued functions of (x, t) with $\Omega(x, 0) = 0 = \alpha(x, 0)$ for every position x of the atom, and $\alpha(x, t)W(x, t)$ is the generalization of the energy operator of the pump field. As an example of $\Omega(x, t)$, we often adopt $\Omega(x, t) = \Omega_0(t)\gamma(x)$ in this paper. Here $\Omega_0(t)$ is a continuous, real-valued function of time $t \geq 0$ with $\Omega_0(0) = 0$, and $\gamma(x)$ a bounded, continuous, real-valued function of the position x of the atom. For instance, $\gamma(x) = \cos kx$.

In the case where $\Omega(x, t) \equiv 0$ and $\alpha(z, t) \equiv 0$, we denote eigenvalues of $H(0, 0; d) := H(\Omega = 0, \alpha = 0; d)$ by $\mathcal{E}_0 < \mathcal{E}_1 < \dots < \mathcal{E}_n < \dots$. When either $\Omega(x, t)$ or $\alpha(t)$ is alive, we denote eigenvalues of $H(\Omega, \alpha; d)$ by $\mathcal{E}_n(\Omega, \alpha; d)$ for $n = 0, 1, \dots$. If the interaction $H_{\text{int}} := i\Omega(t, x)(\sigma_{01}a^{\dagger d} - \sigma_{10}a^d) + \alpha(t)W(x, t)$ is a small perturbation for $H(0, 0; d)$, then each eigenvalue $\mathcal{E}_n(\Omega, \alpha; d)$ sits near its original position $\mathcal{E}_n$, so that the primary order among eigenvalues is kept: $\mathcal{E}_0(\Omega, \alpha; d) < \mathcal{E}_1(\Omega, \alpha; d) < \dots < \mathcal{E}_n(\Omega, \alpha; d) < \dots$. On the other hand, the phase transition of the superradiance [19, 22, 23] tells us about a possibility that $\mathcal{E}_1(\Omega, \alpha; d)$ is less than $\mathcal{E}_0(\Omega, \alpha; d)$ and thus becomes a new ground state energy provided that the interaction H_{int} has some strong strength. For our Hamiltonian, we can classify crossings into two types. One type is the crossing between an ascending eigenvalue $\mathcal{E}_n^+(\Omega, \alpha; d)$ and a descending one $\mathcal{E}_n^-(\Omega, \alpha; d)$ as the strength of the interaction H_{int} grows enough. Another type is the crossing only among descending eigenvalues $\mathcal{E}_n^-(\Omega, \alpha; d)$ (or ascending eigenvalues $\mathcal{E}_n^+(\Omega, \alpha; d)$). We call the former type a *trivial* crossing, and the latter type a *non-trivial* crossing. For the non-trivial crossing, as the strength of the interaction becomes much stronger, even many $\mathcal{E}_n^-(\Omega, \alpha; d)$ s may be less than $\mathcal{E}_0^-(\Omega, \alpha; d)$. We call such a non-trivial crossing the Dicke-type (energy level) crossing [31]. Moreover, $\mathcal{E}_n^-(\Omega, \alpha; d)$ s are capable of usurping the position of the ground state energy in turn. We call such a new ground state energy the *superradiant ground state energy*. The Dicke-type energy level crossings and the appearance of the superradiant ground state energy can be used, together with cavity decay, for carrying away the energy from the system. Based on this idea, we construct mathematical foundations of the concept of superradiant cooling different from that proposed in Ref.15 in and after the next section.

III. THE DICKE-TYPE ENERGY LEVEL CROSSINGS IN THE CASE $\alpha \equiv 0$.

In this section we show how the Dicke-type energy level crossing takes place for $H(\Omega, 0; d)$, that is, for $H(\Omega, \alpha; d)$ in the case where $\alpha(t) \equiv 0$. As in Ref.12 using the well-known identification so that the ground state $|0\rangle$ with the energy ε_0 and the 1st excited state $|1\rangle$ with the energy ε_1 are unitarily equivalent to $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ and $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ respectively, $H(\Omega, 0; d)$ approximately reads

$$H_0(z, t; d) := \begin{pmatrix} -\Delta_c a^\dagger a + \varepsilon_1 - \Delta & -i\Omega(z, t)a^d \\ i\Omega(z, t)a^{\dagger d} & -\Delta_c a^\dagger a + \varepsilon_0 \end{pmatrix}. \quad (3.1)$$

Here we note $\sigma_{00} = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$, $\sigma_{01} = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$, $\sigma_{10} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$, and $\sigma_{11} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$. In their paper we always assume that

$$\varepsilon_1 - \Delta > \varepsilon_0. \quad (3.2)$$

Therefore, the ground state energy of $H_0(z, 0; d)$ (i.e., $H_0(z, t; d)$ with $\Omega(z, t) \equiv 0$) is always ε_0 .

As shown in Sec.A with the method which is a generalization of that in Refs.22 and 45, since $H_0(z, t; d)$ is basically the Hamiltonian of the Jaynes-Cummings model [39], all the energy levels of $H_0(z, t; d)$ are perfectly determined. They are given by $-\Delta_c n + \varepsilon_0$ for non-negative integer n with $n < d$, and $\Xi_n(d) \pm \Upsilon_n(z, t; d)$ for non-negative integer n with $n \geq d$, where

$$\Xi_n(d) = -\Delta_c n + \frac{1}{2}(\varepsilon_0 + \varepsilon_1 + d\Delta_c - \Delta),$$

and $\Upsilon_n(z, t; d)$ is the generalized Rabi frequency [44]:

$$\begin{aligned} & \Upsilon_n(z, t; d) \\ &= \frac{1}{2} \sqrt{(\varepsilon_1 - \varepsilon_0 + d\Delta_c - \Delta)^2 + 4|\Omega(z, t)|^2 \frac{n!}{(n-d)!}}. \end{aligned} \quad (3.3)$$

Here we note all the energy levels of $H_0(z, 0; d)$ are $-\Delta_c n + \varepsilon_0$ and $-\Delta_c n + \varepsilon_1 - \Delta$, $n = 0, 1, \dots$. Therefore, we can conclude that the energy levels of $H_0(z, t; d)$ are completely given by energies $E_n^0(z, t; d)$ and energies $E_n^\pm(z, t; d)$ of the generalized Jaynes-Cummings doublet [17], continuous functions of (z, t) , for each $n = 0, 1, \dots$: For non-negative integers n with $n < d$

$$E_n^0(z, t; d) = -\Delta_c n + \varepsilon_0.$$

For non-negative integers n with $n \geq d$, on the other hand,

$$E_n^-(z, t; d) = \begin{cases} \Xi_n(d) - \Upsilon_n(z, t; d) & \text{if } \varepsilon_1 - \varepsilon_0 \geq \Delta - d\Delta_c, \\ \Xi_{n+d}(d) - \Upsilon_{n+d}(z, t; d) & \text{if } \varepsilon_1 - \varepsilon_0 < \Delta - d\Delta_c, \end{cases}$$

and

$$E_n^+(z, t; d) = \begin{cases} \Xi_{n+d}(d) + \Upsilon_{n+d}(z, t; d) & \text{if } \varepsilon_1 - \varepsilon_0 \geq \Delta - d\Delta_c, \\ \Xi_n(d) + \Upsilon_n(z, t; d) & \text{if } \varepsilon_1 - \varepsilon_0 < \Delta - d\Delta_c. \end{cases}$$

It follows from these definitions that

$$E_n^-(z, 0; d) = \begin{cases} -\Delta_c n + \varepsilon_0 & \text{if } \varepsilon_1 - \varepsilon_0 \geq \Delta - d\Delta_c, \\ -\Delta_c n + \varepsilon_1 - \Delta & \text{if } \varepsilon_1 - \varepsilon_0 < \Delta - d\Delta_c, \end{cases}$$

and

$$\begin{aligned} E_n^+(z, t; d) &\geq E_n^+(z, 0; d) \\ &= \begin{cases} -\Delta_c n + \varepsilon_1 - \Delta & \text{if } \varepsilon_1 - \varepsilon_0 \geq \Delta - d\Delta_c, \\ -\Delta_c n + \varepsilon_0 & \text{if } \varepsilon_1 - \varepsilon_0 < \Delta - d\Delta_c. \end{cases} \end{aligned}$$

The later inequality means that the candidates of super-radiant ground state energy are only $E_n^-(z, t; d)$ s.

We define two spatiotemporal domains $\mathbb{D}_{mn}^{\text{wc}}(d)$ and $\mathbb{D}_{mn}^{\text{sc}}(d)$ for non-negative integers m and n with $\max\{d, m\} < n$ as

$$\begin{aligned} \mathbb{D}_{mn}^{\text{wc}}(d) &:= \{(z, t) \mid E_m^0(z, t; d) < E_n^-(z, t; d) \text{ if } m < d; \\ &\quad E_m^-(z, t; d) < E_n^-(z, t; d) \text{ if } m \geq d\} \end{aligned}$$

and

$$\begin{aligned} \mathbb{D}_{mn}^{\text{sc}}(d) &:= \{(z, t) \mid E_m^0(z, t; d) > E_n^-(z, t; d) \text{ if } m < d; \\ &\quad E_m^-(z, t; d) > E_n^-(z, t; d) \text{ if } m \geq d\} \end{aligned}$$

respectively.

A. In the case $d = 1$

In this subsection we investigate the behavior of the Dicke-type energy level crossings in the case $d = 1$. To do that, we introduce some positive numbers and some domains of the space-time. Then, we divide the whole space of the space-time into three classes, namely, the weak coupling regime, the strong coupling regime, and the critical regime:

For each natural number n , we define a positive number C_{0n}^0 by

$$C_{0n}^0 := \begin{cases} \Delta_c^2 n - \Delta_c (\varepsilon_1 - \varepsilon_0 + \Delta_c - \Delta) & \text{if } \varepsilon_1 - \varepsilon_0 \geq \Delta - \Delta_c, \\ \Delta_c^2 (n+1) - \Delta_c (\varepsilon_1 - \varepsilon_0 + \Delta_c - \Delta) & \text{if } \varepsilon_1 - \varepsilon_0 < \Delta - \Delta_c. \end{cases}$$

We define three domains $\mathcal{D}_{0n}^{\text{wc}}(1)$, $\mathcal{D}_{0n}^{\text{sc}}(1)$, and $\mathcal{D}_{0n}^0(1)$ of the space-time as follows: the spatiotemporal domain $\mathcal{D}_{0n}^{\text{wc}}(1)$ for the weak coupling regime is given by $\mathcal{D}_{0n}^{\text{wc}}(1) := \{(z, t) \mid |\Omega(z, t)|^2 < C_{0n}^0\}$, and the domain $\mathcal{D}_{0n}^{\text{sc}}(1)$ for the strong coupling regime by

$\mathcal{D}_{0n}^{\text{sc}}(1) := \{(z, t) \mid |\Omega(z, t)|^2 > C_{0n}^0\}$. The domain $\mathcal{D}_{0n}^{\text{cr}}(1)$ for the critical regime is defined by $\mathcal{D}_{0n}^{\text{cr}}(1) := \{(z, t) \mid |\Omega(z, t)|^2 = C_{0n}^0\}$.

The following theorem says that how the Dicke-type energy level crossing takes place is completely determined: *Let us suppose $1 < n$ now. Then, the spatiotemporal domain $\mathcal{D}_{0n}^{\text{wc}}(1)$ is equal to the domain $\mathbb{D}_{0n}^{\text{wc}}(1)$, and the spatiotemporal domain $\mathcal{D}_{0n}^{\text{sc}}(1)$ to the domain $\mathbb{D}_{0n}^{\text{sc}}(1)$, i.e., $\mathcal{D}_{0n}^{\text{wc}}(1) = \mathbb{D}_{0n}^{\text{wc}}(1)$ and $\mathcal{D}_{0n}^{\text{sc}}(1) = \mathbb{D}_{0n}^{\text{sc}}(1)$. Namely, the energy level crossing takes place as*

$$E_0^0(z, t; 1) < E_n^-(z, t; 1) \text{ if and only if } (z, t) \text{ in } \mathcal{D}_{0n}^{\text{wc}}(1), \quad (3.4)$$

$$E_0^0(z, t; 1) = E_n^-(z, t; 1) \text{ if and only if } (z, t) \text{ in } \mathcal{D}_{0n}^{\text{cr}}(1), \quad (3.5)$$

$$E_0^0(z, t; 1) > E_n^-(z, t; 1) \text{ if and only if } (z, t) \text{ in } \mathcal{D}_{0n}^{\text{sc}}(1). \quad (3.6)$$

These inequalities (3.4)–(3.6) guarantee the Dicke-type energy level crossing because $E_0^0(z, t; 1)$ and $E_n^-(z, t; 1)$ are continuous functions of the space-time point (z, t) .

When the above Dicke-type energy level crossing between $E_0^0(z, t; 1)$ and $E_n^-(z, t; 1)$ takes place, there is certainly an energy level crossing between $E_m^-(z, t; 1)$ and $E_n^-(z, t; 1)$ for a natural number m with $1 < m < n$. To show it, we introduce two positive constants and define two domains of the space-time:

For natural numbers m, n with $m < n$, we set positive constants C_{mn}^{wc} and C_{mn}^{sc} as

$$C_{mn}^{\text{wc}} = \begin{cases} \Delta_c^2 \left\{ \frac{m+n}{2} + \sqrt{\left(\frac{m+n}{2}\right)^2 + \frac{K^2}{2\Delta_c^2}} \right\} & \text{if } \varepsilon_1 - \varepsilon_0 \geq \Delta - \Delta_c, \\ \Delta_c^2 \left\{ \frac{m+n+2}{2} + \sqrt{\left(\frac{m+n+2}{2}\right)^2 + \frac{K^2}{2\Delta_c^2}} \right\} & \text{if } \varepsilon_1 - \varepsilon_0 < \Delta - \Delta_c, \end{cases}$$

and

$$C_{mn}^{\text{sc}} = \begin{cases} \Delta_c^2 \left\{ m+n+2 + \sqrt{mn + \frac{K^2}{4\Delta_c^2}} \right\} & \text{if } \varepsilon_1 - \varepsilon_0 \geq \Delta - \Delta_c, \\ \Delta_c^2 \left\{ m+n+2 + 2\sqrt{m+n+mn + \frac{K^2}{4\Delta_c^2}} \right\} & \text{if } \varepsilon_1 - \varepsilon_0 < \Delta - \Delta_c, \end{cases}$$

where $K = \varepsilon_1 - \varepsilon_0 + \Delta_c - \Delta$. We give two spatiotemporal domains $\mathcal{D}_{mn}^{\text{wc}}(1)$ and $\mathcal{D}_{mn}^{\text{sc}}(1)$ for non-negative integers m, n with $m < n$ in the following: The domain $\mathcal{D}_{mn}^{\text{wc}}(1)$ for the weak coupling regime is given by $\mathcal{D}_{mn}^{\text{wc}}(1) = \{(z, t) \mid 0 \leq |\Omega(z, t)|^2 < C_{mn}^{\text{wc}}\}$, and the domain $\mathcal{D}_{mn}^{\text{sc}}(1)$ for the strong coupling regime by $\mathcal{D}_{mn}^{\text{sc}}(1) = \{(z, t) \mid |\Omega(z, t)|^2 > C_{mn}^{\text{sc}}\}$. Then, as far as such m goes, the following theorem gives a sufficient condition so that

the crossing between $E_m^-(z, t; 1)$ and $E_n^-(z, t; 1)$ occurs: *Let $1 < m < n$ now. Then, the spatiotemporal domain $\mathcal{D}_{mn}^{\text{wc}}(1)$ is included in the domain $\mathbb{D}_{mn}^{\text{wc}}(1)$, and the domain $\mathcal{D}_{mn}^{\text{sc}}(1)$ in the domain $\mathbb{D}_{mn}^{\text{sc}}(1)$, i.e., $\mathcal{D}_{mn}^{\text{wc}}(1) \subset \mathbb{D}_{mn}^{\text{wc}}(1)$ and $\mathcal{D}_{mn}^{\text{sc}}(1) \subset \mathbb{D}_{mn}^{\text{sc}}(1)$:*

$$E_m^-(z, t; 1) < E_n^-(z, t; 1) \text{ for } (z, t) \text{ in } \mathcal{D}_{mn}^{\text{wc}}(1), \quad (3.7)$$

$$E_m^-(z, t; 1) > E_n^-(z, t; 1) \text{ for } (z, t) \text{ in } \mathcal{D}_{mn}^{\text{sc}}(1). \quad (3.8)$$

These two theorems will be proved at the end of this subsection. But, before proving them, we concretely see some physical situation that they tell us. We employ $\cos 2\pi z$ as $\gamma(z)$. For the position fixed at $z = 0$, the five energies $E_0^0(0, t; 1)$ and $E_n^-(0, t; 1)$, $n = 1, 2, 3, 4$, are numerically calculated as in Fig.1. In the case where the strength $|\Omega_0(t)|$ is given by $|\Omega_0(t)| = 2\kappa, 4\kappa, 6\kappa, 8\kappa$, the two energies $E_0^0(z, t; 1)$ and $E_1^-(z, t; 1)$ are in Fig.2.




FIG. 1: Dicke-type energy level crossings for $\gamma(z) = \cos 2\pi z$. Energies $E_0^0(0, t; 1)$ (solid line), $E_1^-(0, t; 1)$ (dashed), $E_2^-(0, t; 1)$ (short-dashed), $E_3^-(0, t; 1)$ (dotted), and $E_4^-(0, t; 1)$ (dashed-dotted). The physical parameters are set as $\varepsilon_0 = 0$, $\varepsilon_1 = 6\kappa$, $\Delta = 1\kappa$, $\Delta_c = -3\kappa$ with a unit κ . The position z is fixed at $z = 0$.

Suppose that $\varepsilon_1 - \varepsilon_0 \geq \Delta - \Delta_c$ now. Let our atom be in the state with the energy $E_m^-(z_0, t_0; 1)$ at an initial space-time point (z_0, t_0) in the domain $\mathcal{D}_{mn}^{\text{wc}}(1)$ for a non-negative integer m and a positive integer n with $0 \leq m < n$. The atom is apt to sit in the state with energy $E_m^-(z_0, t_0; 1)$ because being in the state with energy $E_m^-(z_0, t_0; 1)$ is more stable than in the state with energy $E_n^-(z_0, t_0; 1)$. Once, however, a space-time point (z, t) plunges into the domain $\mathcal{D}_{mn}^{\text{sc}}(1)$, the energy level crossing takes place as shown in its process (3.4)–(3.6). Thus, being in the state with energy $E_m^-(z, t; 1)$ is *not* stable any longer, and thus, it goes down to the state with energy $E_n^-(z, t; 1)$. This descent can be caused by cavity decay even if we cannot expect atomic decay as pointed out in Ref.12. The circumstance for such a cooling is usually not in thermal equilibrium, and thus, the temperature which this system loses is not given by thermodynamic temperature. Nevertheless, according to the thermodynamics low as in Eq.(5.1) of Ref.40 and Eq.(2.1) of Ref.41, we roughly estimate the effective temperature $\Delta T_{m \rightarrow n}$




FIG. 2: Dicke-type energy level crossings for $\gamma(z) = \cos 2\pi z$. Energies $E_0^0(z, t; 1)$ (solid line), $E_1^-(z, t; 1)$ with $|\Omega_0(t)| = 2\kappa$ (dashed), $E_1^-(z, t; 1)$ with $|\Omega_0(t)| = 4\kappa$ (short-dashed), $E_1^-(z, t; 1)$ with $|\Omega_0(t)| = 6\kappa$ (dotted), and $E_1^-(z, t; 1)$ with $|\Omega_0(t)| = 8\kappa$ (dashed-dotted). The time is fixed so that $|\Omega_0(t)| = 2\kappa, 4\kappa, 6\kappa, 8\kappa$.

coming from the descent. Then, we can expect that the crossing between $E_m^-(z, t; 1)$ and $E_n^-(z, t; 1)$ makes the temperature go down at most $\Delta T_{m \rightarrow n}$ estimated at:

$$\Delta T_{m \rightarrow n} \approx \frac{2(n-m)}{k_B} \left| \frac{|\Omega(z, t)|^2}{\Upsilon_m(z_0, t_0; 1) + \Upsilon_n(z, t; 1)} - |\Delta_c| \right| + \frac{2m}{k_B} \left| \frac{|\Omega(z, t)|^2 - |\Omega(z_0, t_0)|^2}{\Upsilon_m(z_0, t_0; 1) + \Upsilon_n(z, t; 1)} \right| \quad (3.9)$$

for the point (z_0, t_0) in the spatiotemporal domain $\mathcal{D}_{mn}^{\text{wc}}(1)$ and the point (z, t) in the domain $\mathcal{D}_{mn}^{\text{sc}}(1)$, of course, provided that there is nothing to obstruct the temperature loss. Here k_B is the Boltzmann constant and $\Upsilon_\ell(z, t; 1) = (1/2)\sqrt{K^2 + 4|\Omega(z, t)|^2}^\ell$ the generalized Rabi frequency (3.3) for each natural number ℓ .

Conversely, let our atom be in the state with the energy $E_n^-(z_0, t_0; 1)$ at an initial space-time point (z_0, t_0) in the domain $\mathcal{D}_{mn}^{\text{sc}}(1)$. Then, since a process reverse to the above energy level crossing takes place if the coupling regime recoils from the spatiotemporal domain $\mathcal{D}_{mn}^{\text{sc}}(1)$ to the domain $\mathcal{D}_{mn}^{\text{wc}}(1)$, this reverse crossing can also carry away the temperature $\Delta T_{n \rightarrow m}$ of the atom so that $\Delta T_{n \rightarrow m} = \Delta T_{m \rightarrow n}$ for the point (z_0, t_0) in the spatiotemporal domain $\mathcal{D}_{mn}^{\text{sc}}(1)$ and the point (z, t) in the domain $\mathcal{D}_{mn}^{\text{wc}}(1)$. We illustrate this situation for $E_1^-(z, t; 1)$ and $E_3^-(z, t; 1)$ in Fig.3. The diagrammatic illustration about temperature loss is represented by the thick arrows in Fig.3. Therefore, our arguments say that there is a possibility of the following mechanism for superradiant cooling: Let a space-time point $(z_{2\ell+1}, t_{2\ell+1})$ be in the domain $\mathcal{D}_{01}^{\text{wc}}(1)$ and a space-time point $(z_{2\ell+2}, t_{2\ell+2})$ in the domain $\mathcal{D}_{01}^{\text{sc}}(1)$, respectively, for each $\ell = 0, 1, \dots, N-1$ with a natural number N . The Dicke-type energy level crossings and their reverse crossings may carry away the




FIG. 3: Energy decay between $E_1^-(z, t; 1)$ (solid line) and $E_3^-(z, t; 1)$ (dashed). The physical parameters are set as in Fig.1 and the strength $|\Omega_0(t)|$ is fixed at 12κ . $\gamma(z) = \cos 2\pi z$ in $\Omega(z, t)$. The thick arrows stand for the temperature loss.

temperature of the atom:

$$\frac{2}{k_B} \sum_{\nu=1}^{2N} \frac{|\Omega(z_\nu, t_\nu)|^2}{|K|/2 + \Upsilon_1(z_\nu, t_\nu; 1)} - \frac{4N}{k_B} |\Delta_c| \quad (3.10)$$

at most by Eq.(3.9) if $K \geq 0$. We can make similar argument when $K < 0$.

When the whole space of the space-time is $\mathbb{D}_{01}^{\text{wc}}(1)$, we find the Sisyphus-type mechanism in the energy spectrum as the group of Ritsch pointed out in Ref.12. To see the diagrammatic representation we consider a concrete example now. For the strength $|\Omega_0(t)| = 2\kappa$ we have the cavity-induced atom cooling as in Fig.4. Up-arrows in




FIG. 4: Cavity-induced atom cooling (bold line) in $\mathbb{D}_{01}^{\text{wc}}(1)$. Energies $E_0^0(z, t; 1)$ (solid line) and $E_0^-(z, t; 1)$ (dashed). The physical parameters are set as in Fig.1.

Fig.4 represent the energy that the atom coupled with photons gains by photon absorption, namely, we have to throw a driving laser to the atom for its excitement. Down-arrows mean the energy loss caused by cavity decay. In this domain $\mathbb{D}_{01}^{\text{wc}}(1)$ of the space-time we cannot find such a sequence $\{(z_\nu, t_\nu)\}_{\nu=1}^{2L}$ of the space-time points. Thus, Eq.(3.10) does not work. As shown in the

following concrete example, on the other hand, we can expect the sequence $\{(z_\nu, t_\nu)\}_{\nu=1}^{2N}$: If we take a strength as $|\Omega_0| = 8\kappa$, then Fig.4 changes to Fig.5, and then, we can find the sequence $\{(z_\nu, t_\nu)\}_{\nu=1}^{2N}$ in Fig.5. Compare




FIG. 5: Dicke-type crossings and cavity-induced atom cooling (bold line). Energies $E_0^0(z, t; 1)$ (solid line) and $E_1^-(z, t; 1)$ (dashed). The physical parameters are set as in Fig.1.

Fig.4 and Fig.5. In Fig.5 we can make only down-arrows without any uparrow. It means that the system loses energy only because of cavity decay. Namely, we do not have to throw the driving laser to the atom in the cavity for its excitement. We note this type emission comes from a kind of superradiance [19, 22, 23]. Thus, this energy-spectral property may give a mechanism for another superradiant cooling for a two-level atom in the strong coupling regime, as well as self-organized, cooperative atoms [15, 16]. As we can realize it from Figs.2 and 5, the energy loss by cavity decay gets large as the strength $|\Omega_0(t)|$ of the coupling grows large. We give the surfaces of $E_0^0(z, t; 1)$ and $E_1^-(z, t; 1)$ as functions of (z, t) in Figs.6, 7, and 8.




FIG. 6: Surfaces of energies $E_0^0(z, t; 1) = 0$ (dashed line) and $E_1^-(z, t; 1)$ (solid). The physical parameters are set as in Fig.1.




FIG. 7: Surface of energy $E_1^-(z, t; 1)$ (solid line), where $E_0^0(z, t; 1) = 0$. The physical parameters are set as in Fig.1.




FIG. 8: Surface of energy $E_1^-(z, t; 1)$ (solid line), where $E_0^0(z, t; 1) = 0$. The physical parameters are set as in Fig.1.

We prove our two theorems on the Dicke-type energy level crossing now: To make the calculations below simple, we set K as $K := \varepsilon_1 - \varepsilon_0 + \Delta_c - \Delta$ again.

We give a proof of crossings (3.4)–(3.6) first. Let us suppose $K \geq 0$ now. It is easy to show the equation,

$$E_0^0(z, t; 1) - E_n^-(z, t; 1) = \Delta_c n - \frac{1}{2}K + \Upsilon_n(z, t; 1), \quad (3.11)$$

where $\Upsilon_n(z, t; 1)$ is the generalized Rabi frequency (3.3). Let the symbol $\#$ denote either $>$, $=$, or $<$. After multiplying both sides of the expression $|\Omega(z, t)|^2 \# \Delta_c^2 n - \Delta_c K$ by $4n$, add the term K^2 to both sides of the multiplied expression. Then, we know that the expression $|\Omega(z, t)|^2 \# \Delta_c^2 n - \Delta_c K$, is equivalent to the expression $K^2 + 4n\Omega(z, t)^2 \# 4(\Delta_c^2 n^2 - \Delta_c K n + K^2/4)$. Since $-\Delta_c n + K/2 \geq 0$, we can take the square root of the last expression, and thus, it is equivalent to $2\Upsilon_n(z, t; 1) \# 2(-\Delta_c n + K/2)$. Therefore, Eq.(3.11) says

that

$$E_0^0(z, t; 1) - E_n^-(z, t; 1) \neq 0 \iff |\Omega(z, t)|^2 \neq \Delta_c^2 n - \Delta_c K, \quad (3.12)$$

where “RHS” $\iff$ “LHS” means that the right hand side of the equation is *equivalent* to the left hand side. The equivalence (3.12) brings the crossings (3.4)–(3.6).

Let us suppose $K < 0$ now. Then, in the same way we did in the case $K > 0$, we have the equivalence:

$$\begin{aligned} E_0^0(z, t; 1) - E_n^-(z, t; 1) \neq 0 \\ \iff \Upsilon_{n+1}(z, t; 1) \neq -\Delta_c(n+1) + \frac{1}{2}K. \end{aligned} \quad (3.13)$$

The term $-\Delta_c(n+1) + K/2$ is positive because we assumed the condition (3.2) and $\Delta_c < 0$. Thus, by taking the square of both sides of the right expression of the equivalence (3.13) and following the way we did in the case $K \geq 0$, we know the expression $E_0^0(z, t; 1) - E_n^-(z, t; 1) \neq 0$ is equivalent to the expression $|\Omega(z, t)|^2 \neq \Delta_c^2(n+1) - \Delta_c K$, which implies our desired result.

We consider the proofs of the crossing (3.7) and (3.8) next. Let us suppose $d \leq m < n$. A direct calculation leads to

$$\begin{aligned} E_m^-(z, t; 1) - E_n^-(z, t; 1) \\ = \Delta_c(n-m) + \Upsilon_n(z, t; 1) - \Upsilon_m(z, t; 1) \\ = (n-m) \left[\Delta_c + \frac{|\Omega(z, t)|^2}{\Upsilon_n(z, t; 1) + \Upsilon_m(z, t; 1)} \right]. \end{aligned} \quad (3.14)$$

Since $n > m$ and $|\Delta_c| = -\Delta_c$, Eq.(3.14) says that the expression $E_m^-(z, t; 1) - E_n^-(z, t; 1) \neq 0$ is equivalent to the expression $|\Omega(z, t)|^2 (\Upsilon_n(z, t; 1) + \Upsilon_m(z, t; 1))^{-1} \neq |\Delta_c|$. Multiplying both sides of this by $2|\Delta_c|^{-1}|\Omega(z, t)|^{-1}(\Upsilon_n(z, t; 1) + \Upsilon_m(z, t; 1))$, we realize that the later expression is equivalent to the expression $2|\Omega(z, t)|^{-1}/|\Delta_c| \neq \sqrt{(K/|\Omega(z, t)|)^2 + 4n} + \sqrt{(K/|\Omega(z, t)|)^2 + 4m}$. Hence it follows from these equivalences that

$$E_m^-(z, t; 1) - E_n^-(z, t; 1) \neq 0 \iff 2 \frac{|\Omega(z, t)|}{|\Delta_c|} - \sqrt{\left(\frac{K}{|\Omega(z, t)|}\right)^2 + 4m} \neq \sqrt{\left(\frac{K}{|\Omega(z, t)|}\right)^2 + 4n}. \quad (3.15)$$

Suppose $K \geq 0$ now. Here we note a simple inequality for non-negative numbers A, B , and C : $(\sqrt{A+B} + \sqrt{A+C})^2 \geq A+B+A+C$. Using this, we can show that the expression $4(|\Omega(z, t)|/|\Delta_c|)^2 < (K/|\Omega(z, t)|)^2 + 4n + (K/|\Omega(z, t)|)^2 + 4m$ implies the expression $4(|\Omega(z, t)|/|\Delta_c|)^2 < \{\sqrt{(K/|\Omega(z, t)|)^2 + 4n} + \sqrt{(K/|\Omega(z, t)|)^2 + 4m}\}^2$.

Taking the square root of both sides of this inequality implies the expression, $2(|\Omega(z, t)|/|\Delta_c|) < \sqrt{(K/|\Omega(z, t)|)^2 + 4n} + \sqrt{(K/|\Omega(z, t)|)^2 + 4m}$. Therefore, we obtain the following:

$$\begin{aligned} 4 \left(\frac{|\Omega(z, t)|}{|\Delta_c|} \right)^2 &< \left(\frac{K}{|\Omega(z, t)|} \right)^2 + 4n + \left(\frac{K}{|\Omega(z, t)|} \right)^2 + 4m \\ \implies 2 \frac{|\Omega(z, t)|}{|\Delta_c|} - \sqrt{\left(\frac{K}{|\Omega(z, t)|}\right)^2 + 4m} &< \sqrt{\left(\frac{K}{|\Omega(z, t)|}\right)^2 + 4n}, \end{aligned} \quad (3.16)$$

where “RHS” $\implies$ “LHS” means that the right hand side of the equation *implies* the left hand side.

We define a polynomial $g_{wc}(r)$ by $g_{wc}(r) := 2r^2 - 2(m+n)r - K^2/|\Delta_c|^2$. Multiplying both sides of this by 2, the inequality $g_{wc}(|\Omega(z, t)|/|\Delta_c|) < 0$ is equivalent to the

inequality $4(|\Omega(z, t)|/|\Delta_c|)^4 < 4(m+n)(|\Omega(z, t)|/|\Delta_c|)^2 + 2(K/|\Delta_c|)^2$. Multiplying both sides of this newly obtained inequality by $(|\Delta_c|/|\Omega(z, t)|)^2$ we reach the following:

$$\begin{aligned}
& g_{\text{wc}} \left(\left(\frac{|\Omega(z, t)|}{|\Delta_c|} \right)^2 \right) < 0 \\
\iff & 4 \left(\frac{|\Omega(z, t)|}{|\Delta_c|} \right)^2 < \left(\frac{K}{|\Omega(z, t)|} \right)^2 + 4m + \left(\frac{K}{|\Omega(z, t)|} \right)^2 + 4n.
\end{aligned} \tag{3.17}$$

It follows from the implication (3.16), and equivalences (3.15) and (3.17) that

$$\begin{aligned}
& g_{\text{wc}} \left(\left(\frac{|\Omega(z, t)|}{|\Delta_c|} \right)^2 \right) < 0 \\
\implies & E_m^-(z, t; 1) - E_n^-(z, t; 1) < 0.
\end{aligned} \tag{3.18}$$

Since the point r_0^{wc} defined by $r_0^{\text{wc}} := (m+n)/2 + \sqrt{((m+n)/2)^2 + K^2/(2\Delta_c^2)}$ satisfies $g_{\text{wc}}(r_0^{\text{wc}}) = 0$, we have the inequality $g_{\text{wc}}(r) < g_{\text{wc}}(r_0^{\text{wc}}) = 0$ provided that $0 \leq r < r_0^{\text{wc}}$. This fact tells us that

$$\begin{aligned}
& \left(\frac{|\Omega(z, t)|}{|\Delta_c|} \right)^2 < \frac{m+n}{2} + \sqrt{\left(\frac{m+n}{2} \right)^2 + \frac{K^2}{2\Delta_c^2}} \\
\implies & g_{\text{wc}} \left(\left(\frac{|\Omega(z, t)|}{|\Delta_c|} \right)^2 \right) < 0.
\end{aligned} \tag{3.19}$$

Therefore, we can conclude the inequality (3.7) from implications (3.18) and (3.19).

We prove the inequality (3.8) next. Multiply both sides of the inequality $(|\Omega(z, t)|/|\Delta_c|)^2 - (n-m) > (|\Omega(z, t)|/|\Delta_c|)\sqrt{(K/|\Omega(z, t)|)^2 + 4m}$ by 4, and add $(K/|\Omega(z, t)|)^2$ to both sides of the multiplied inequality. Then, we know that the inequality $(|\Omega(z, t)|/|\Delta_c|)^2 - (n-m) > (|\Omega(z, t)|/|\Delta_c|)\sqrt{(K/|\Omega(z, t)|)^2 + 4m}$ implies the inequality $4(|\Omega(z, t)|/|\Delta_c|)^2 + (K/|\Omega(z, t)|)^2 - 4(n-m) > (K/|\Omega(z, t)|)^2 + 4(|\Omega(z, t)|/|\Delta_c|)\sqrt{(K/|\Omega(z, t)|)^2 + 4m}$, which is equivalent to the inequality

$$\begin{aligned}
& 4 \left(\frac{|\Omega(z, t)|}{|\Delta_c|} \right)^2 - 4 \left(\frac{|\Omega(z, t)|}{|\Delta_c|} \right) \sqrt{\left(\frac{K}{|\Omega(z, t)|} \right)^2 + 4m} \\
& \quad + \left\{ \left(\frac{K}{|\Omega(z, t)|} \right)^2 + 4m \right\} \\
& > \left(\frac{K}{|\Omega(z, t)|} \right)^2 + 4n.
\end{aligned}$$

Taking the square root of both sides of the last inequality, we reach the implication:

$$\begin{aligned}
& \left(\frac{|\Omega(z, t)|}{|\Delta_c|} \right)^2 - (n-m) > \left(\frac{|\Omega(z, t)|}{|\Delta_c|} \right) \sqrt{\left(\frac{K}{|\Omega(z, t)|} \right)^2 + 4m} \\
\implies & 2 \frac{|\Omega(z, t)|}{|\Delta_c|} - \sqrt{\left(\frac{K}{|\Omega(z, t)|} \right)^2 + 4m} > \sqrt{\left(\frac{K}{|\Omega(z, t)|} \right)^2 + 4n}.
\end{aligned} \tag{3.20}$$

We define a polynomial $g_{\text{sc}}(r)$ by $g_{\text{sc}}(r) := r^2 - 2(m+n)r + (n-m)^2 - K^2/|\Delta_c|^2$ this time. We note the inequality $g_{\text{sc}}(|\Omega(z, t)|/|\Delta_c|)^2 > 0$ is equivalent to the inequality $(|\Omega(z, t)|/|\Delta_c|)^4 - 2(n-m)(|\Omega(z, t)|/|\Delta_c|)^2 + (n-m)^2 > (|\Omega(z, t)|/|\Delta_c|)^2 \{ (K/|\Omega(z, t)|)^2 + 4m \}$. Here

we added $4m(|\Omega(z, t)|/|\Delta_c|)^2 + (K/|\Delta_c|)^2$ to both sides of $g_{\text{sc}}(|\Omega(z, t)|/|\Delta_c|)^2 > 0$, and then, the right hand side was factorized with the factor $(|\Omega(z, t)|/|\Delta_c|)^2$. Thus, taking the square root of both side of this inequality, we obtain the following implication:

$$\begin{aligned}
& g_{\text{sc}} \left(\left(\frac{|\Omega(z, t)|}{|\Delta_c|} \right)^2 \right) > 0 \\
\implies & \left(\frac{|\Omega(z, t)|}{|\Delta_c|} \right)^2 - (n-m) > \frac{|\Omega(z, t)|}{|\Delta_c|} \sqrt{\left(\frac{K}{|\Delta_c|} \right)^2 + 4m}.
\end{aligned} \tag{3.21}$$

It follows from the implications (3.20) and (3.21), and equivalence (3.15) that

$$g_{\text{sc}} \left(\left(\frac{|\Omega(z, t)|}{|\Delta_c|} \right)^2 \right) > 0 \\ \implies E_m^-(z, t; 1) - E_n^-(z, t; 1) > 0. \quad (3.22)$$

Since $r_0^{\text{sc}} := m + n + \sqrt{4mn + K^2/\Delta_c^2}$ satisfies $g_{\text{sc}}(r_0^{\text{sc}}) = 0$, we have the inequality $g_{\text{sc}}(r) > g_{\text{sc}}(r_0^{\text{sc}}) = 0$ provided that $r_0^{\text{sc}} < r$. This fact tells us that

$$\left(\frac{|\Omega(z, t)|}{|\Delta_c|} \right)^2 > m + n + 2\sqrt{mn + \frac{K^2}{4\Delta_c^2}} \\ \implies g_{\text{sc}} \left(\left(\frac{|\Omega(z, t)|}{|\Delta_c|} \right)^2 \right) > 0. \quad (3.23)$$

Therefore, we can conclude the inequality (3.8) from implications (3.22) and (3.23).

In the case where $K < 0$, we have

$$E_m^-(z, t; 1) - E_n^-(z, t; 1) \\ = (n - m) \left[\Delta_c + \frac{\Omega(z, t)^2}{\Upsilon_{n+1}(z, t; 1) + \Upsilon_{m+1}(z, t; 1)} \right].$$

Hence it follows from this that we obtain the crossing (3.7) and (3.8) by using $m + 1$ and $n + 1$ instead of m and n respectively in the above argument for the case where $K \geq 0$.

B. In the case $d \geq 2$

We assume $d \geq 2$ in this subsection. Let m and n be non-negative integers satisfying $m < d \leq n$. We set a positive number $C_{mn}^0(d)$ by

$$C_{mn}^0(d) := \begin{cases} \frac{(n-d)!}{n!} (n-m) |\Delta_c| \left\{ |\Delta_c| (n-m) + K_d \right\} & \text{if } \varepsilon_1 - \varepsilon_0 \geq \Delta - d\Delta_c, \\ \frac{n!}{(n+d)!} (n+d-m) |\Delta_c| \times & \\ \quad \times \left\{ |\Delta_c| (n+d-m) + K_d \right\} & \text{if } \varepsilon_1 - \varepsilon_0 < \Delta - d\Delta_c, \end{cases}$$

where $K_d := \varepsilon_1 - \varepsilon_0 + d\Delta_c - \Delta$. Then, we define three domains $\mathcal{D}_{mn}^{\text{wc}}(d)$, $\mathcal{D}_{mn}^{\text{sc}}(d)$, and $\mathcal{D}_{mn}^{\text{cr}}(d)$ of the space-time in the following: the domain $\mathcal{D}_{mn}^{\text{wc}}(d)$ for the weak coupling regime is given by $\mathcal{D}_{mn}^{\text{wc}}(d) := \{(z, t) \mid |\Omega(z, t)|^2 < C_{mn}^0(d)\}$, and the domain $\mathcal{D}_{mn}^{\text{sc}}(d)$ for the strong coupling regime by $\mathcal{D}_{mn}^{\text{sc}}(d) := \{(z, t) \mid |\Omega(z, t)|^2 > C_{mn}^0(d)\}$. The domain $\mathcal{D}_{mn}^{\text{cr}}(d)$ for the critical regime is defined by $\mathcal{D}_{mn}^{\text{cr}}(d) := \{(z, t) \mid |\Omega(z, t)|^2 = C_{mn}^0(d)\}$.

In the case $d \geq 2$ we can show the following theorem: the domain $\mathbb{D}_{mn}^{\text{wc}}(d)$ is equal to the domain $\mathcal{D}_{mn}^{\text{wc}}(d)$, and the domain $\mathbb{D}_{mn}^{\text{sc}}(d)$ to the domain $\mathcal{D}_{mn}^{\text{sc}}(d)$, i.e.,

$\mathbb{D}_{mn}^{\text{wc}}(d) = \mathcal{D}_{mn}^{\text{wc}}(d)$ and $\mathbb{D}_{mn}^{\text{sc}}(d) = \mathcal{D}_{mn}^{\text{sc}}(d)$, for every m and n with $m < d \leq n$. Namely, the energy level crossing takes place as

$$E_m^0(z, t; d) < E_n^-(z, t; d) \text{ if and only if } (z, t) \text{ in } \mathcal{D}_{mn}^{\text{wc}}(d), \quad (3.24)$$

$$E_m^0(z, t; d) = E_n^-(z, t; d) \text{ if and only if } (z, t) \text{ in } \mathcal{D}_{mn}^{\text{cr}}(d), \quad (3.25)$$

$$E_m^0(z, t; d) > E_n^-(z, t; d) \text{ if and only if } (z, t) \text{ in } \mathcal{D}_{mn}^{\text{sc}}(d). \quad (3.26)$$

Here we note

$$C_{0n}^0(d) = \Delta_c^2 \frac{n}{(n-1) \cdots (n-d+1)} \\ + |\Delta_c| \frac{K_d}{(n-1) \cdots (n-d+1)} \\ < \Delta_c^2 n + |\Delta_c| K \leq C_{0n}^0$$

because $d\Delta_c < \Delta_c < 0$. Thus, it follows from this inequality that $\mathcal{D}_{0n}^{\text{sc}}(1) \subset \mathcal{D}_{0n}^{\text{sc}}(d)$. It means that *non-linear coupling* (i.e., $d \geq 2$) causes the Dicke-type energy level crossing easier than linear coupling (i.e., $d = 1$) does.

Before proving our desired statements, we set K_d as $K_d := \varepsilon_1 - \varepsilon_0 + d\Delta_c - \Delta$ for the simplicity in calculation. It is easy to check that

$$E_m^0(z, t; d) - E_n^-(z, t; d) \\ = \begin{cases} -\Delta_c(m-n) - \frac{K_d}{2} + \Upsilon_n(z, t; d) & \text{if } K_d \geq 0, \\ -\Delta_c(m-n-d) - \frac{K_d}{2} + \Upsilon_{n+d}(z, t; d) & \text{if } K_d < 0, \end{cases}$$

which implies the following equivalence:

$$E_m^0(z, t; d) - E_n^-(z, t; d) \neq 0 \\ \iff \begin{cases} \Upsilon_n(z, t; d) \neq -\Delta_c(n-m) + \frac{K_d}{2} & \text{if } K_d \geq 0, \\ \Upsilon_{n+d}(z, t; d) \neq -\Delta_c(n+d-m) + \frac{K_d}{2} & \text{if } K_d < 0. \end{cases} \quad (3.27)$$

We see the energy level crossing in the case where $K_d \geq 0$ first. Thus, suppose $K_d \geq 0$. Since $|\Delta_c| = -\Delta_c$, multiplying both sides of the expression $|\Omega(z, t)|^2 \neq C_{mn}^0(d)$ by $n!/(n-d)!$, we know that the expression is equivalent to the expression $|\Omega(z, t)|^2 n!/(n-d)! \neq \Delta_c^2 (n-m)^2 - \Delta_c(n-m)K_d$. Adding $K_d^2/4$ to both sides of the multiplied expression and factorizing the left hand side with $1/4$, we know the expression is equivalent to the expression $(K_d^2 + 4|\Omega(z, t)|^2 n!/(n-d)!)/4 \neq \Delta_c^2 (n-m)^2 - \Delta_c(n-m)K_d + K_d^2/4$. Taking the square root of both sides of this expression, we obtain the equivalence:

$$|\Omega(z, t)|^2 \neq C_{mn}^0(d) \\ \iff \Upsilon_n(z, t; 1) \neq -\Delta_c(n-m) + \frac{K_d}{2} \quad (3.28)$$

since $\Delta_c < 0$, $m < n$, and $0 \leq K_d$. Finally, by the equivalences (3.27) and (3.28), we reach the equivalence:

$$E_m^0(z, t; d) - E_n^-(z, t; d) \# 0 \iff |\Omega(z, t)|^2 \# C_{mn}^0(d), \quad (3.29)$$

which says that the desired energy level crossing takes place when $K_d \geq 0$.

We see the energy level crossing in the case where $K_d < 0$ next. Let us suppose $K_d < 0$ now. Note the inequalities $-\Delta_c(n + d - m) + K_d/2 \geq (\varepsilon_1 - \varepsilon_0 - d\Delta_c - \Delta)/2 > (\varepsilon_1 - \varepsilon_0 - \Delta_c - \Delta)/2 > (\varepsilon_1 - \varepsilon_0 - \Delta)/2 > 0$ because of the assumption (3.2) and the condition $-\Delta_c > 0$. Thus, in the same way we had the equivalence (3.28), we obtain the equivalence,

$$|\Omega(z, t)|^2 \# C_{mn}^0(d) \iff \Upsilon_{n+d}(z, t; d) \# -\Delta_c(n + d - m) + \frac{K_d}{2}. \quad (3.30)$$

Then, the equivalences (3.27) and (3.30) bring the equivalence (3.29), which secures our statement in the case $K_d < 0$.

In the case where $d \geq 2$, there is not always a ground state of $H_0(z, t; d)$. Namely, there is a case that the minimum energy of $H_0(z, t; d)$ does not exist. To see this fact, we assume $\varepsilon_1 - \varepsilon_0 \approx \Delta - d\Delta_c$ for simplicity. Then, since we have

$$\begin{aligned} E_n^-(z, t; d) - E_{n+1}^-(z, t; d) &= \Delta_c + |\Omega(z, t)| \sqrt{n(n-1) \cdots (n-d+2)} \times \\ &\quad \times \left\{ \sqrt{n+1} - \sqrt{n-d+1} \right\} \\ &> \Delta_c + |\Omega(z, t)| \sqrt{n(n-1) \cdots (n-d+2)} \times \\ &\quad \times \left\{ \sqrt{n+1} - \sqrt{n-1} \right\} \\ &= \Delta_c + 2|\Omega(z, t)| \frac{\sqrt{(n-1) \cdots (n-d+2)}}{\sqrt{1+1/n} + \sqrt{1-1/n}} \\ &> \Delta_c + |\Omega(z, t)| \sqrt{(d-1)!}, \end{aligned}$$

where we used the inequalities, $n > d \geq 2$ and $\sqrt{1+1/n} + \sqrt{1-1/n} \geq 2$. The last inequality says that the Hamiltonian $H_0(z, t; d)$ does not have a ground state for (z, t) satisfying $|\Omega(z, t)| > |\Delta_c|/\sqrt{(d-1)!}$ because

$$\cdots < E_{n+1}^-(z, t; d) < E_n^-(z, t; d) < \cdots < E_{d+1}^-(z, t; d)$$

in the case where $d \geq 2$.

IV. SUPERRADIANT GROUND STATE ENERGY IN THE CASE $\alpha \equiv 0$.

As we knew at the end of the previous section, once the Dicke-type crossing takes place in the case $d \geq 2$, there is *every* possibility that $H_0(z, t; d)$ is not bounded from below. Thus, to make sure of the existence of the superradiant ground state energy, we consider only the case $d = 1$ in this section.

Set a positive number Θ_n for $n = 1, 2, \dots$ as

$$\Theta_n = \begin{cases} 2n\Delta_c^2 + |\Delta_c|\sqrt{4n^2\Delta_c^2 + K^2} & \text{if } \varepsilon_1 - \varepsilon_0 \geq \Delta - \Delta_c, \\ 2(n+1)\Delta_c^2 + |\Delta_c|\sqrt{4(n+1)^2\Delta_c^2 + K^2} & \text{if } \varepsilon_1 - \varepsilon_0 < \Delta - \Delta_c, \end{cases}$$

where $K = \varepsilon_1 - \varepsilon_0 + \Delta_c - \Delta$. Using this number, we define a domain $\mathcal{G}_n(1)$ of the time-space by

$$\mathcal{G}_n(1) = \left\{ (z, t) \mid \sqrt{\Theta_n} < |\Omega(z, t)| < \sqrt{\Theta_{n+1}} \right\}$$

for each $n = 1, 2, \dots$. We can prove the following theorem: (i) when a space-time point (z, t) is in $\mathcal{G}_n(1)$, the point (z, t) is always in $\mathcal{D}_{0n}^{\text{sc}}(1)$. Namely, the Dicke-type energy level crossing takes place for that point (z, t) in $\mathcal{G}_n(1)$; (ii) the superradiant ground state energy $\inf \text{Spec}(H_0(z, t; 1))$ appears as:

$$\inf \text{Spec}(H_0(z, t; 1)) = \min\{E_n^-(z, t; 1), E_{n+1}^-(z, t; 1)\} \quad (4.1)$$

for (z, t) in $\mathcal{G}_n(1)$.

To prove our theorem we set K as $K := \varepsilon_1 - \varepsilon_0 + \Delta_c - \Delta$ again. Part (i) follows from the following easy inequalities: $C_{0n}^0 < 2n\Delta_c^2 + |\Delta_c|K \leq \Theta_n$ if $K \geq 0$; $C_{0n}^0 < 2(n+1)\Delta_c^2 + |\Delta_c|K \leq \Theta_n$ if $K < 0$. For a non-negative number r , define a function $g(r)$ by $g(r) := |\Delta_c|r + L - (1/2)\sqrt{K^2 + 4|\Omega(z, t)|^2r}$, where $L = (\varepsilon_0 + \varepsilon_1 + \Delta_c - \Delta)$. Then, for every positive number r the expression $g'(r) \# 0$ is equivalent to the expression $\Delta_c^2(K^2 + 4|\Omega(z, t)|^2r) \# |\Omega(z, t)|^4$. Hence it follows from this that

$$g'(r) \# 0 \iff r \# r_0 := \frac{|\Omega(z, t)|^2}{4\Delta_c^2} - \frac{K^2}{4|\Omega(z, t)|^2}, \quad (4.2)$$

provided that $r_0 > 0$. It is evident that the number r_0 is positive if and only if the inequality $|\Omega(z, t)|^2 > |\Delta_c K|$ holds. By the equivalence (4.2) together with this fact, we have the implication:

$$|\Omega(z, t)|^2 > |\Delta_c K| \implies \inf_{r \geq 0} g(r) = g(r_0). \quad (4.3)$$

Simple calculation leads to the equivalence:

$$\begin{aligned} n &\leq \frac{|\Omega(z, t)|^2}{4\Delta_c^2} - \frac{K^2}{4|\Omega(z, t)|^2} < n+1 \\ \iff \begin{cases} 0 \leq |\Omega(z, t)|^4 - 4n\Delta_c^2|\Omega(z, t)|^2 - \Delta_c^2K^2, \\ |\Omega(z, t)|^4 - 4(n+1)\Delta_c^2|\Omega(z, t)|^2 - \Delta_c^2K^2 < 0. \end{cases} \end{aligned} \quad (4.4)$$

We define two functions $g_j(r)$, $j = 1, 2$, for positive number r by $g_1(r) := r^2 - 4n\Delta_c^2r - \Delta_c^2K^2$ and $g_2(r) := r^2 - 4(n+1)\Delta_c^2r - \Delta_c^2K^2$. Then, setting r_{j0} as $r_{10} := 2n\Delta_c^2 + |\Delta_c|\sqrt{K^2 + 4n^2\Delta_c^2}$ and $r_{20} := 2(n+1)\Delta_c^2 + |\Delta_c|\sqrt{K^2 + 4(n+1)^2\Delta_c^2}$ respectively, we have

$g_1(r) > g_1(r_{10}) = 0$ if $r > r_{10}$ and $g_2(r) < g_2(r_{20}) = 0$ if $0 < r < r_{20}$. Set a non-negative number θ_n for each $n = 1, 2, \dots$ as $\theta_n := 2n\Delta_c^2 + |\Delta_c|\sqrt{K^2 + 4n^2\Delta_c^2}$. Then, we have $\theta_n \geq |\Delta_c K|$. So, inserting $|\Omega(z, t)|^2$ into r of the above inequalities, we reach the implication:

$$\begin{aligned} \theta_n &< |\Omega(z, t)|^2 < \theta_{n+1} \\ \implies \text{RHS of the equivalence (4.4) is satisfied.} \end{aligned} \quad (4.5)$$

Combining the results (4.3), (4.4), and (4.5), we obtain the implication:

$$\begin{aligned} \theta_n &< |\Omega(z, t)|^2 < \theta_{n+1} \\ \implies g(n) \text{ or } g(n+1) \text{ is located nearest } \inf_{r \geq 0} g(r). \end{aligned} \quad (4.6)$$

Noting $E_n^-(z, t; 1) = g(n)$ if $K \geq 0$; $E_n^-(z, t; 1) = g(n+1)$ if $K < 0$, and $\Theta_n = \theta_n$ if $K \geq 0$; $\Theta_n = \theta_{n+1}$ if $K < 0$, we obtain part (ii).

V. STABILITY OF DICKE-TYPE CROSSINGS IN THE CASE $\alpha \neq 0$.

In this section we consider the case where $d = 1$ too. we show the stability of the Dicke-type energy level crossings under the effect of the generalized energy operator $\alpha(t)W(z, t)$ of the pump field for sufficiently small strength $|\alpha(t)|$. Our $W(z, t)$ includes $i\alpha(a - a^\dagger)$, of course. To show this fact we employ the method for proving Theorem 4.3(v) of Ref.31. In the same way we did in Sec.III, through the well-known identification as in Ref.12, $H(\Omega, \alpha; 1)$ reads $H_\alpha(z, t; 1) := H_0(z, t; 1) + \alpha(z, t)W(z, t)$. We recall $\alpha(z, 0) = 0$ for all the position z .

For the Hilbert space representing the state space in which $H_\alpha(z, t; d)$ acts, we denote the inner product of the Hilbert space by $\langle \Psi | \Phi \rangle$.

For every positive number ϵ , we denote $C_\epsilon(z, t)$ by

$$\begin{aligned} C_\epsilon(z, t) \\ := (1 + \epsilon) \left\{ 1 + \left(\frac{\epsilon}{1 + \epsilon} \right)^2 (\epsilon_0 + |\epsilon_1 - \Delta|) \right. \\ \left. + \left(\frac{1 + \epsilon}{\epsilon} \right)^2 \frac{|\Omega(z, t)|^2}{4\Delta_c^2} \right\}^{1/2}. \end{aligned}$$

Set a positive number $C_{0n}^0[\theta]$ for each natural number n and a non-negative number θ as:

$$\begin{aligned} C_{0n}^0[\theta] \\ := \begin{cases} (\theta - \Delta_c)^2 n + (\theta - \Delta_c)(\epsilon_1 - \epsilon_0 + \Delta_c - \Delta) & \text{if } \epsilon_1 - \epsilon_0 \geq \Delta - \Delta_c, \\ (\theta - \Delta_c)^2 (n + 1) + (\theta - \Delta_c)(\epsilon_1 - \epsilon_0 + \Delta_c - \Delta) & \text{if } \epsilon_1 - \epsilon_0 < \Delta - \Delta_c, \end{cases} \end{aligned}$$

and set a positive number $C[n]$ for each natural number n as:

$$\begin{aligned} C[n] \\ := \begin{cases} \Delta_c^2 n + \frac{\epsilon_0(\epsilon_1 + \Delta_c - \Delta)}{n} & \text{if } \epsilon_1 - \epsilon_0 \geq \Delta - \Delta_c, \\ \Delta_c^2 (n + 1) + \frac{\epsilon_0(\epsilon_1 + \Delta_c - \Delta)}{n + 1} & \text{if } \epsilon_1 - \epsilon_0 < \Delta - \Delta_c. \end{cases} \end{aligned}$$

Moreover, for every positive function $f(z, t)$, and positive numbers b and ϵ' , we define a domain $\mathcal{D}(\epsilon, \epsilon'; b, f)$ of the space-time by

$$\begin{aligned} \mathcal{D}(\epsilon, \epsilon'; b, f) \\ := \left\{ (z, t) \mid |\alpha(z, t)| < \frac{\epsilon'}{2b} \text{ and } |\alpha(z, t)| \left(bC_\epsilon(z, t) + f(z, t) \right) < \frac{\epsilon'(1 + \epsilon)}{2} \right\}. \end{aligned}$$

Also the domain $\mathcal{D}_{0n}^{\text{sc}}(1; \theta)$ of the space-time is defined by

$$\begin{aligned} \mathcal{D}_{0n}^{\text{sc}}(1; \theta) \\ := \{ (z, t) \mid |\Omega(z, t)|^2 > C_{0n}^0[\theta] \text{ and } |\Omega(z, t)|^2 > C[n] \}. \end{aligned}$$

We prove the following theorem concerning the stability of the Dicke-type energy level crossings: *Let our $W(z, t)$ satisfy the conditions (A1)–(A3):*

(A1) *For every space-time point (z, t) , $W(z, t)$ is a symmetric operator so that it can act the all states on which $H_0(z, 0; 1)$ acts;*

(A2) *there is a positive constant b_1 so that*

$$\begin{aligned} \langle W(z, t)\Psi | W(z, t)\Psi \rangle^{1/2} \\ \leq b_1 \langle H_0(z, 0; 1)\Psi | H_0(z, 0; 1)\Psi \rangle^{1/2} + b_2(z, t) \langle \Psi | \Psi \rangle^{1/2} \end{aligned} \quad (5.1)$$

for all states Ψ on which $H_0(z, 0; 1)$ acts and every space-time point (z, t) , where $b_2(z, t)$ is some positive function of (z, t) ;

(A3) *there is a constant ϵ with $0 < \epsilon < 1$ so that $|\alpha(z, t)| < \{b_1(1 + \epsilon)\}^{-1}$ for all the space-time points (z, t) .*

Then, the Hamiltonian $H_\alpha(z, t; 1)$ has an eigenvalue $\mathcal{E}_n^{\mathfrak{h}}(z, t; 1)$ near $E_n^{\mathfrak{h}}(z, t; 1)$ for each non-negative integer n , where $\mathfrak{h} = 0, \pm$. Moreover, there is a constant κ_0 with $0 < \kappa_0 < 1/4$ so that the Dicke-type energy level crossing takes place between the eigenvalues $\mathcal{E}_0^0(z, t; 1)$ and $\mathcal{E}_n^-(z, t; 1)$ in the process from the space-time point $(z, 0)$ to the space-time point (z_, t_*) , provided that the*

latter point (z_*, t_*) is in $\mathcal{D}(\epsilon, \kappa_0; b_1, b_2) \cap \mathcal{D}_{0n}(1; \theta)$ for a number θ with $\theta > 0$. Therefore, $H_\alpha(z_*, t_*; 1)$ has the superradiant ground state energy.

We prove this theorem below. Set H_I as $H_I := H_0(z, t; 1) - H_0(z, 0; 1)$ for simplicity. The symbol H_0 stands for $H_0(z, 0; 1)$ from now on. Here we recall $\Omega(z, 0) = 0$ for every position z .

In the same way that we proved Lemma 4.2 of Ref.31, we can estimate $\langle W(z, t)\Psi | W(z, t)\Psi \rangle$ from above by using $\langle H_0(z, t; d)\Psi | H_0(z, t; d)\Psi \rangle$ and $\langle \Psi | \Psi \rangle$. Using the canonical commutation relation $[a, a^\dagger] = 1$ leads to the equation:

$$\langle H_I \Psi | H_I \Psi \rangle = |\Omega(z, t)|^2 (\langle \Psi | a^\dagger a | \Psi \rangle + \langle \sigma_{11} \Psi | \sigma_{11} \Psi \rangle),$$

which easily implies the inequality

$$\langle H_I \Psi | H_I \Psi \rangle \leq |\Omega(z, t)|^2 (\langle \Psi | a^\dagger a | \Psi \rangle + \langle \Psi | \Psi \rangle).$$

By using the Schwarz inequality and the inequality $XY \leq (\eta X + Y/4\eta)^2$ for every $X, Y \geq 0$ and arbitrary $\eta > 0$, we obtain the inequality:

$$\langle \Psi | a^\dagger a | \Psi \rangle \leq \left(\eta \langle a^\dagger a \Psi | a^\dagger a \Psi \rangle^{1/2} + \langle \Psi | \Psi \rangle^{1/2} / (4\eta) \right)$$

for arbitrary $\eta > 0$. Simple calculation yields

$$\begin{aligned} & \langle H_0 \Psi | H_0 \Psi \rangle \\ &= \Delta_c^2 \langle a^\dagger a \Psi | a^\dagger a \Psi \rangle + (\varepsilon_1 - \Delta) \langle \sigma_{11} \Psi | \sigma_{11} \Psi \rangle \\ & \quad + \varepsilon_0 \langle \sigma_{00} \Psi | \sigma_{00} \Psi \rangle, \end{aligned}$$

so that

$$\begin{aligned} & \langle a^\dagger a \Psi | a^\dagger a \Psi \rangle \\ &= (1/\Delta_c^2) \left\{ \langle H_0 \Psi | H_0 \Psi \rangle - (\varepsilon_1 - \Delta) \langle \sigma_{11} \Psi | \sigma_{11} \Psi \rangle \right. \\ & \quad \left. - \varepsilon_0 \langle \sigma_{00} \Psi | \sigma_{00} \Psi \rangle \right\}. \end{aligned}$$

Combining these inequalities and setting η as $\eta := |\Delta_c| \delta / (\sqrt{2} |\Omega(z, t)|)$ for arbitrary $\delta > 0$, we reach the inequality:

$$\begin{aligned} & \langle H_I \Psi | H_I \Psi \rangle^{1/2} \\ & \leq |\Omega(z, t)| \left\{ 2\eta^2 \langle a^\dagger a \Psi | a^\dagger a \Psi \rangle + \left(\frac{1}{8\eta^2} + 1 \right) \langle \Psi | \Psi \rangle \right\}^{1/2} \\ & \leq \left\{ \delta^2 \langle H_0 \Psi | H_0 \Psi \rangle + \Theta(\delta; \Omega)^2 \langle \Psi | \Psi \rangle \right\}^{1/2} \\ & \leq \delta \langle H_0 \Psi | H_0 \Psi \rangle^{1/2} + \Theta(\delta; \Omega) \langle \Psi | \Psi \rangle^{1/2}, \end{aligned} \quad (5.2)$$

where

$$\Theta(\delta; \Omega) = \sqrt{1 + \delta^2(\varepsilon_0 + |\varepsilon_1 - \Delta|) + \frac{|\Omega(z, t)|^2}{4\Delta_c^2 \delta^2}}.$$

Applying the triangle inequality $\langle (A + B)\Psi | (A + B)\Psi \rangle^{1/2} \leq \langle A\Psi | A\Psi \rangle^{1/2} + \langle B\Psi | B\Psi \rangle^{1/2}$ to the term

$\langle (H_0(z, t; 1) - H_I)\Psi | (H_0(z, t; 1) - H_I)\Psi \rangle^{1/2}$, we have the inequality:

$$\begin{aligned} \langle H_0 \Psi | H_0 \Psi \rangle^{1/2} & \leq \langle H_0(z, t; 1)\Psi | H_0(z, t; 1)\Psi \rangle^{1/2} \\ & \quad + \langle H_I \Psi | H_I \Psi \rangle^{1/2}. \end{aligned} \quad (5.3)$$

Inequalities (5.2) and (5.3) tell us that the term $\langle H_0 \Psi | H_0 \Psi \rangle^{1/2}$ is bounded from above as:

$$\begin{aligned} \langle H_0 \Psi | H_0 \Psi \rangle^{1/2} & \leq \langle H_0(z, t; 1)\Psi | H_0(z, t; 1)\Psi \rangle^{1/2} \\ & \quad + \delta \langle H_0 \Psi | H_0 \Psi \rangle^{1/2} + \Theta(\delta; \Omega) \langle \Psi | \Psi \rangle^{1/2}, \end{aligned}$$

which implies the inequality,

$$\begin{aligned} (1 - \delta) \langle H_0 \Psi | H_0 \Psi \rangle^{1/2} & \leq \langle H_0(z, t; 1)\Psi | H_0(z, t; 1)\Psi \rangle^{1/2} \\ & \quad + \Theta(\delta; \Omega) \langle \Psi | \Psi \rangle^{1/2}. \end{aligned} \quad (5.4)$$

Set δ as $0 < \delta := \epsilon(1 + \epsilon)^{-1} < 1$ for arbitrary number ϵ with $0 < \epsilon < 1$ now. Then, combining the inequalities (5.1) and (5.4) we can conclude that

$$\begin{aligned} & \langle W(z, t)\Psi | W(z, t)\Psi \rangle^{1/2} \\ & \leq b_1(1 + \epsilon) \langle H_0(z, t; 1)\Psi | H_0(z, t; 1)\Psi \rangle^{1/2} \\ & \quad + (b_1 C_\epsilon(z, t) + b_2(z, t)) \langle \Psi | \Psi \rangle^{1/2}. \end{aligned} \quad (5.5)$$

Thus, applying Lemma 4.1 of Ref.31 and Theorem 6.29 in III §6 of Ref.42 to our Hamiltonian $H_\alpha(z, t; 1) = H_0 + H_I$, we know that the Hamiltonian $H_\alpha(z, t; 1)$ has an eigenvalue $\mathcal{E}_n^a(z, t; 1)$ for each $n = 0, 1, 2, \dots$.

Define an operator $T(\kappa)$ as the closure of $H_0(z, t; 1) + \kappa W(z, t)$ for every complex number κ . By applying Theorem 2.6 and Remark 2.7 in VII §2 of Ref.42 to the inequality (5.5), the operator $T(\kappa)$ is an analytic family of type (A) for every $\kappa \in \mathbb{C}$ with $|\kappa| < \{b_1(1 + \epsilon)\}^{-1}$. Thus, taking $\alpha(z, t)$ as this κ , i.e., $H_\alpha(z, t; 1) = T(\alpha(z, t))$, Theorem 3.9 in VII §3 of Ref.42 says that $\mathcal{E}_n^a(z, t; 1)$ is a continuous function of (z, t) by the assumption (A3), and it sits near $E_n^a(z, t; 1)$. So, as shown in Sec.III, candidates which makes the Dicke-type energy level crossing are $E_n^-(z, t; 1)$'s. Namely, for $H_\alpha(z, t; 1)$, candidates which makes the Dicke-type energy level crossing are $\mathcal{E}_n^-(z, t; 1)$'s.

Let E_n and $\mathcal{E}_n(\kappa)$ be respectively eigenvalues of $H_0(z, t; 1)$ and $T(\kappa)$ satisfying $\mathcal{E}_n(0) = E_n$. Following the regular perturbation theory (see §XII of Ref.43), the eigenvalue $\mathcal{E}_n(\kappa)$ has the expression,

$$\begin{aligned} \mathcal{E}_n(\kappa) &= \frac{\langle \Phi_n | T(\kappa) P_n(\kappa) \Phi_n \rangle}{\langle \Phi_n | P_n(\kappa) \Phi_n \rangle} \\ &= E_n + \kappa \frac{\langle \Phi_n | W(z, t) P_n(\kappa) \Phi_n \rangle}{\langle \Phi_n | P_n(\kappa) \Phi_n \rangle} \end{aligned} \quad (5.6)$$

where $P_n(\kappa)$ is the orthogonal projection given by

$$P_n(\kappa) = -\frac{1}{2\pi i} \oint_{|E - E_n| = \tilde{\epsilon}} (T(\kappa) - E)^{-1} dE$$

with a sufficiently small $\tilde{\epsilon} > 0$, and Φ_n is a normalized eigenvector of $H_0(z, t; 1)$ satisfying $H_0(z, t; 1)\Phi_n = E_n\Phi_n$.

Since $W(z, t)$ is symmetric, use of the Schwarz inequality makes the inequality:

$$\left| \frac{\langle \Phi_n | W(z, t) P_n(\kappa) \Phi_n \rangle}{\langle \Phi_n | P_n(\kappa) \Phi_n \rangle} \right| \leq \frac{\langle W(z, t) \Phi_n | W(z, t) \Phi_n \rangle^{1/2}}{\langle P_n(\kappa) \Phi_n | P_n(\kappa) \Phi_n \rangle^{1/2}}. \quad (5.7)$$

To get the right hand side of the above inequality, we used the equations $P_n(\kappa)^2 = P_n(\kappa) = P_n(\kappa)^*$ in the denominator, and the inequality $\langle P_n(\kappa) \Phi_n | P_n(\kappa) \Phi_n \rangle \leq 1$ in the numerator. By the inequality (5.5) we can estimate $\langle W(z, t) \Phi_n | W(z, t) \Phi_n \rangle^{1/2}$ as $\langle W(z, t) \Phi_n | W(z, t) \Phi_n \rangle^{1/2} \leq b_1(1 + \epsilon)E_n + b_1C_\epsilon(z, t) + b_2(z, t)$. Here we note that $0 \leq \langle P_n(\kappa) \Phi_n | P_n(\kappa) \Phi_n \rangle \rightarrow 1$ as $|\kappa| \rightarrow 0$, so that we have $1/2 < \langle P_n(\kappa) \Phi_n | P_n(\kappa) \Phi_n \rangle$ for sufficiently small $|\kappa|$. Thus, combining these with the inequality (5.7), there is a positive constant κ_0 so that

$$|\mathcal{E}_n(\kappa) - E_n| \leq 2|\kappa| \{b_1(1 + \epsilon)E_n + b_1C_\epsilon(z, t) + b_2(z, t)\} \quad (5.8)$$

if $|\kappa| \leq \kappa_0$.

Now we take the coupling strength $|\alpha(z, t)|$ for space-time point $(z, t) \in \mathcal{D}(\epsilon, \kappa_0; b_1, b_2)$ as the coupling parameter κ . Let us define a positive number κ_1 as $\kappa_1 := \kappa_0(1 + \epsilon)$. Then, it is easy to check that $2|\alpha(z, t)|b_1(1 + \epsilon) < \kappa_1$ and that $2|\alpha(z, t)|\{b_1C_\epsilon(z, t) + b_2(z, t)\} < \kappa_1$. Combining these inequalities with the inequality (5.8), we obtain the inequality:

$$(1 - \kappa_1)E_n - \kappa_1 \leq \mathcal{E}_n(\alpha(z, t)) \leq (1 + \kappa_1)E_n + \kappa_1 \quad (5.9)$$

for space-time point $(z, t) \in \mathcal{D}(\epsilon, \kappa_0; b_1, b_2)$. From now on, we take the κ_0 and ϵ so that $0 < \kappa_1 < 1/2$.

Let us set a number L as $L := \varepsilon_0 + \varepsilon_1 + \Delta_c - \Delta$ again. It is easy to show that $\Omega_n(z, t; 1)^2 - \Xi_n(1)^2 = -\Delta_c^2 n^2 + (|\Omega(z, t)|^2 + \Delta_c^2 L)n - \varepsilon_0(\varepsilon_1 + \Delta_c - \Delta)$, and thus, we obtain the equivalence:

$$\begin{aligned} \Omega_n(z, t; 1) = |\Omega_n(z, t; 1)| &> |\Xi_n(1)| \geq \Xi_n(1) \\ \iff |\Omega(z, t)|^2 &> \Delta_c^2 n - \Delta_c L + \frac{\varepsilon_0(\varepsilon_1 + \Delta_c - \Delta)}{n}. \end{aligned}$$

Hence it follows from this that $E_n^-(z, t; 1)$ is negative for the point $(z, t) \in \mathcal{D}_{0n}^{\text{sc}}(1; \theta)$.

In the same way we did to get the equivalence (3.12), we obtain the equivalence in the following. In the case $K \equiv \varepsilon_1 - \varepsilon_0 + \Delta_c - \Delta$ is non-negative, for every $\theta \geq 0$ and each natural number n we have

$$\begin{aligned} E_0^0(z, t; 1) \nmid E_n^-(z, t; 1) + n\theta \\ \iff |\Omega(z, t)|^2 \nmid (\theta - \Delta_c)^2 n + (\theta - \Delta_c)K \equiv C_{0n}^0[\theta]. \end{aligned} \quad (5.10)$$

Let K be negative now. We note the inequality $(\theta - \Delta_c)\{(\theta - \Delta_c)(n + 1) + K\} \geq (\theta - \Delta_c)\{-\Delta_c + K\} \geq 0$ because of the condition (3.2). Thus, in the same way

we did to get the equivalence (3.12), for every $\theta \geq 0$ and each natural number n we have

$$\begin{aligned} E_0^0(z, t; 1) \nmid E_n^-(z, t; 1) + (n + 1)\theta \\ \iff |\Omega(z, t)|^2 \nmid (\theta - \Delta_c)^2(n + 1) + (\theta - \Delta_c)K \equiv C_{0n}^0[\theta]. \end{aligned} \quad (5.11)$$

Thus, we obtain that $E_0^0(z, t; 1) > (1 + \theta)E_n^-(z, t; 1) + \theta$ since $1 < 1 + \theta$ and $E_n^-(z, t; 1) < 0$ for every point $(z, t) \in \mathcal{D}_{0n}^{\text{sc}}(1; \theta)$. We take the θ defined by the equation $\kappa_1 = \theta/(2 + \theta)$ now. Then, the following inequality holds: $E_0^0(z, t; 1) > (1 + \kappa_1)E_n^-(z, t; 1)/(1 - \kappa_1) + 2\kappa_1/(1 - \kappa_1)$, which implies

$$(1 - \kappa_1)E_0^0(z, t; 1) - \kappa_1 > (1 + \kappa_1)E_n^-(z, t; 1) + \kappa_1 \quad (5.12)$$

for every $(z, t) \in \mathcal{D}_{0n}^{\text{sc}}(1; \theta)$.

Combining the inequalities (5.9) and (5.12) leads to the inequality:

$$\begin{aligned} \mathcal{E}_n^-(z, t; 1) &\leq (1 + \kappa_1)E_n^-(z, t; 1) + \kappa_1 \\ &< (1 - \kappa_1)E_0^0(z, t; 1) - \kappa_1 < \mathcal{E}_0^0(z, t; 1) \end{aligned} \quad (5.13)$$

for every $(z, t) \in \mathcal{D}(\epsilon, \kappa_0; b_1, b_2) \cap \mathcal{D}_{0n}^{\text{sc}}(1; \theta)$.

Since $\mathcal{E}_n^0(z, t; 1)$ is a continuous function of (z, t) and $\mathcal{E}_0^0(z, 0; 1) = E_0^0(z, 0; 1) < E_n^-(z, 0; 1) = \mathcal{E}_n^-(z, 0; 1)$, the inequality (5.13) means that the Dicke-type energy level crossing takes place.

VI. CONCLUSION

We have showed that the system of a two-level atom coupled with a laser in a cavity has the Dicke-type energy level crossing in the process that the atom-cavity interaction of the system undergoes changes between the weak coupling regime and the strong one. By using the Dicke-type energy level crossing, we have found the following two possibilities in (mathematical) theory for the cavity-induced atom cooling. We can use a laser only for controlling the strength of the atom-cavity interaction without throwing another laser to the atom for driving it to the excited state, and moreover, we can obtain much larger energy loss caused by cavity decay, if we obtain the cavity that implements the domain $\mathcal{D}_{0n}^{\text{sc}}(1)$ of the space-time. Based on these results, we can say that we lay mathematical foundations for the concept of another superradiant cooling in addition to that proposed by Domokos and Ritsch. Adding the mathematical foundations to the idea of the cavity-induced atom cooling by Ritsch *et al.*, we can also say that the process of our superradiant cooling requires only cavity decay and control of the position of the atom, without atomic absorption and emission of photons. Therefore, whether the mechanism of our superradiant cooling can primarily be demonstrated or not depends on whether we can make such a fine cavity that the spatiotemporal domain $\mathcal{D}_{01}^{\text{sc}}(1)$ for the strong coupling regime can be implemented or not [6, 7, 8, 9, 46, 47, 48, 49, 50, 51, 52].

Acknowledgments

This work is partially supported by Japan Society for the Promotion of Science (JSPS), Grand-in-Aid for Scientific Research (C) 20540171.

APPENDIX A: THE EIGENVALUE PROBLEM FOR $H_0(z, t; d)$

In this appendix, to solve the eigenvalue problem: $H_0(z, t; 1)\Psi(g) = E\Psi(g)$, we adopt the way that we did in §3 and §6 of Ref.31 into our calculations. Set $\lambda := i\Omega(z, t)$ for simplicity.

Let $n < d$ for a while. For a complex constant g set $\Psi(g) := \begin{pmatrix} 0 \\ ga^\dagger{}^n\psi_0 \end{pmatrix}$, where ψ_0 is the vacuum state of the photon field of our laser. It follows immediately that the condition, $H(d; \Omega, 0)\Psi(g) = E\Psi(g)$, is equivalent to the condition, $E = \omega n$.

Let $n \geq d$ now. Set $\Psi(g) := \begin{pmatrix} a^\dagger{}^{n-d}\psi_0 \\ ga^\dagger{}^n\psi_0 \end{pmatrix}$ this time. In

the same way as in Ref.31, we conclude that the condition, $H(d; \Omega, 0)\Psi(g) = E\Psi(g)$, is equivalent to the conditions,

$$\begin{cases} (n-d)\omega + \lambda^* \binom{n}{d} d!g = E - \mu, \\ \lambda + ng\omega = Eg. \end{cases}$$

Solving these equations, we obtain

$$g = \frac{d\omega - \mu \pm \sqrt{(\mu - d\omega)^2 + 4 \binom{n}{d} d!|\lambda|^2}}{2\lambda^* \binom{n}{d} d!}.$$

and

$$E = n\omega + \frac{\mu - d\omega}{2} \pm \frac{1}{2} \sqrt{(\mu - d\omega)^2 + 4 \binom{n}{d} d!\lambda^2}.$$

-
- [1] W. Neuhauser, M. Hohenstatt, P. E. Toschek, and H. Dehmelt, Phys. Rev. A **22**, 1137 (1980).
 - [2] S. Chu, L. W. Hollberg, J. E. Bjorkholm, A. Cable, and A. Ashkin, Phys. Rev. Lett. **55**, 48 (1985).
 - [3] C. Cohen-Tannoudji and W. D. Phillips, Physics Today **43**, 33 (1990).
 - [4] E. A. Cornell and C. E. Wieman, Rev. Mod. Phys. **74**, 875 (2002).
 - [5] W. Ketterle, Rev. Mod. Phys. **74**, 1131 (2002).
 - [6] C. J. Hood, M. S. Chapman, T. W. Lynn, and H. J. Kimble, Phys. Rev. Lett. **80**, 4157 (1998).
 - [7] J. M. Raimond, M. Brune, and S. Haroche, Rev. Mod. Phys. **75**, 565 (2001).
 - [8] H. Mabuchi and A. C. Doherty, Science **298**, 1372 (2002).
 - [9] J. McKeever, A. Boca, A. D. Boozer, J. R. Buck, and H. J. Kimble, Nature **425**, 268 (2003).
 - [10] S. M. Dutra, *Cavity Quantum Electrodynamics* (Wiley-Interscience Publication, New York 2005).
 - [11] S. Haroche and J.-M. Raimond, *Exploring the Quantum: Atoms, Cavities, and Photons* (Oxford University Press, Oxford, 2006).
 - [12] P. Horak, G. Hechenblaikner, K. M. Gheri, H. Stecher, and H. Ritsch, Phys. Rev. Lett. **79**, 4974 (1997).
 - [13] C. J. Hood, T. W. Lynn, A. C. Doherty, A. S. Parkins, and H. J. Kimble, Science **287**, 1447 (2000).
 - [14] T. Fischer, P. Maunz, P. W. H. Pinkse, T. Puppe, and G. Rempe, Phys. Rev. Lett. **88**, 163002 (2002).
 - [15] P. Domokos and H. Ritsch, J. Opt. Soc. Am. B **20**, 1098 (2003).
 - [16] P. Domokos and H. Ritsch, Phys. Rev. Lett. **25**, 253003 (2002).
 - [17] C. C. Gerry and P. L. Knight, *Introductory Quantum Optics* (Cambridge University Press, Cambridge, 2005).
 - [18] A. V. Andreev, V. I. Emel'yanov, and Yu. A. Il'inskii, *Cooperative Effects in Optics* (Institute of Physics Publishing, Bristol, 1993).
 - [19] J.-S. Peng and G. X. Li, *Introduction to Modern Quantum Optics* (World Scientific, Singapore, 1998).
 - [20] G. Preparata, *QED Coherence in Matter* (World Scientific, Singapore, 1995).
 - [21] C. P. Enz, Helv. Phys. Acta **70**, 141 (1997).
 - [22] M. Hirokawa, Rev. Math. Phys. **13**, 221 (2001).
 - [23] M. Hirokawa, Phys. Lett. A **294**, 13 (2002).
 - [24] J. Stenger, S. Inouye, D. M. Stamper-Kurn, A. P. Chikkatur, D. E. Pritchard, and W. Ketterle, Appl. Phys. B **69**, 347 (1999).
 - [25] S. Inouye, A. P. Chikkatur, D. M. Stamper-Kurn, J. Stenger, D. E. Pritchard, and W. Ketterle, Science **285**, 571 (1999).
 - [26] D. Schneble, Y. Torii, M. Boyd, E. W. Streed, D. E. Pritchard, and W. Ketterle, Science **300**, 475 (2003).
 - [27] L. Fallani, C. Fort, N. Piovella, M. Cola, F. S. Cataliotti, M. Inguscio, and R. Bonifacia, Phys. Rev. A **71**, 033612 (2005).
 - [28] J. V. Pulé, A. F. Verbeure, and V. A. Zagrebnev, J. Phys. A: Math. Gen. **38**, 5173 (2005).
 - [29] J. V. Pulé, A. F. Verbeure, and V. A. Zagrebnev, J. Phys. A: Math. Gen. **37**, L321 (2004).
 - [30] J. V. Pulé, A. F. Verbeure, and V. A. Zagrebnev, J. Stat. Phys. **119**, 309 (2005).
 - [31] M. Hirokawa, to appear in Indiana Univ. Math. J.
 - [32] R. H. Dicke, Phys. Rev. **93**, 99 (1954).
 - [33] A. Parmeggiani and M. Wakayama, Forum Math. **14**, 539 (2002); *ibid.* **14**, 669 (2002); *ibid.* **15**, 955 (2003).
 - [34] K. Nagatou, M. T. Nakao, and M. Wakayama, Numer. Funct. Anal. Optim. **23**, 633 (2003).
 - [35] A. Parmeggiani, Kyushu J. Math. **58**, 277 (2004); Comm. Math. Phys. **279**, 285 (2008); *Introduction to the spectral theory of non-commutative harmonic oscillators* (COE Lecture Note vol.8, Kyushu University, The 21st Century COE Program "DMHF", Fukuoka, 2008).
 - [36] T. Ichinose and M. Wakayama, Commun. Math. Phys. **256**, 697 (2005); Rep. Math. Phys. **59**, 421 (2007).

- [37] K. Hepp and E. H. Lieb, Ann. Phys. (N.Y.) **76**, 360 (1973); Phys. Rev. A **8**, 2517 (1973); Helv. Phys. Acta **46** 573 (1973).
- [38] M. O. Scully and M. S. Zubairy, *Quantum Optics* (Cambridge Univ. Press, 2006).
- [39] E. T. Jaynes and F. W. Cummings, Proc. IEEE **51**, 89 (1963).
- [40] H. J. Metcalf and P. van der Straten, *Laser Cooling and Trapping* (Springer-Verlag, New York, 1999).
- [41] F. Bardou, J.-P. Bouchaud, A. Aspect, and C. Cohen-Tannoudji, *Lévy Statistics and Laser Cooling* (Cambridge University Press, Cambridge, 2002).
- [42] T. Kato, *Perturbation theory for linear operators* (Springer-Verlag, New York, 1995).
- [43] M. Reed and B. Simon, *Methods of modern Mathematical Physics IV. Analysis of Operators* (Academic Press, San Diego, 1978).
- [44] P. W. Milonni and J. H. Eberly, *Lasers* (Wiley Interscience Publication, New York, 1988).
- [45] S. M. Barnett and P. M. Radmore, *Methods in Theoretical Quantum Optics* (Oxford University Press, Oxford, 2002).
- [46] R. J. Thompson, G. Rempe, and H. J. Kimble, Phys. Rev. Lett. **68**, 1132 (1992).
- [47] M. Brune, P. Nussenzweig, F. Schmidt-Kaler, F. Bernardot, A. Maali, J. M. Raimond, and S. Haroche, Phys. Rev. Lett. **72**, 3339 (1994).
- [48] M. Brune, F. Schmidt-Kaler, A. Maali, J. Dreyer, E. Hagley, J. M. Raimond, and S. Haroche, Phys. Rev. Lett. **76**, 1800 (1996).
- [49] J. J. Childs, K. An, M. S. Otteson, R. R. Dasari, and M. S. Feld, Phys. Rev. Lett. **77**, 2901 (1996).
- [50] P. Grangier, Science **281**, 56 (1998).
- [51] H. Nha, Y.-T. Chough, and K. An, J. Korean Phys. Soc. **37**, 693 (2000).
- [52] W. T. M. Irvine, K. Hennessy, and D. Bouwmeester, Phys. Rev. Lett. **96**, 057405 (2006).